

# The Resolved Kennicutt-Schmidt Law in Nearby Galaxies

R. Momose<sup>1,2,3</sup>, J. Koda<sup>4</sup>, R. C. Kennicutt, Jr<sup>5</sup>, F. Egusa<sup>6</sup>,  
S. K. Okumura<sup>3,7</sup>, D. Calzetti<sup>8</sup>, G. Liu<sup>8,9</sup>, J. Donovan Meyer<sup>4</sup>,  
N. Z. Scoville<sup>10</sup>, T. Sawada<sup>11</sup>, N. Kuno<sup>12</sup>

<sup>1</sup>ICRR, University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba, 277-8582, Japan;  
email: [momo.s.rieko@nao.ac.jp](mailto:momo.s.rieko@nao.ac.jp);

<sup>2</sup>University of Tokyo; <sup>3</sup>NAOJ; <sup>4</sup>SUNY; <sup>5</sup>IoA; University of Cambridge, <sup>6</sup>JAXA; <sup>7</sup>Japan Women's University; <sup>8</sup>UMASS; <sup>9</sup>JHU; <sup>10</sup>Caltech; <sup>11</sup>Joint ALMA Office; <sup>12</sup>NRO, NAOJ

**Abstract.** The Kennicutt-Schmidt law (Schmidt 1959; Kennicutt 1998, hereafter K-S law) is a power law correlation between area averaged star formation rate ( $\Sigma_{SFR}$ ) and gas surface density ( $\Sigma_{gas}$ ). Despite its importance, the physics that underlie this correlation has remained unclear. The power law index,  $N$ , is a prime discriminator of the mechanisms that regulate star formation and form the K-S law (e.g. Leroy *et al.* 2008; Tan 2010). We present a study of the resolved K-S law for 10 nearby disk galaxies using our new CO(1-0) data at 750 and 500 pc resolutions. The CO(1-0) line emission is established as a tracer of the molecular gas column density, and results in a super-linear correlation ( $N = 1.3$  and  $1.8$ ). We discuss the cause of the discrepancy between previous studies, and the mechanism of star formation indicated from our new results.

**Keywords.** stars: formation — galaxies: spiral — ISM: molecules

## 1. Results and Discussions

We derive the K-S law on 750 pc scale by employing a similar procedure as Bigiel *et al.* (2008, hereafter B08), but with CO(1-0). The index at this scale shows a super-linear correlation with  $N=1.3\pm0.02$ . Furthermore we verify the K-S law on a 500 pc scale. The diffuse emission from both H $\alpha$  and 24  $\mu$ m subtraction is also examined on this scale as an example. The best fit linear regressions also result in a super-linear slope both without and with diffuse emission subtraction ( $N=1.2\pm0.05$  and  $N=1.8\pm0.09$ , respectively).

Our result is different from B08 ( $N \sim 1$ ). We find that the discrepancy of the slope between our results and B08 is caused largely by the adopted molecular gas tracer, CO(1-0) or CO(2-1). CO(2-1) has only a factor of 2-3 higher  $n_{crit}$  ( $n_{crit} \geq 10^3 \text{ cm}^{-3}$ ) and level energy (equivalent temperature of 16.5 K); and thus, is excited more when the density and/or temperature is higher than the typical (a few  $100 \text{ cm}^{-3}$ ; 10 K) even just mildly. In fact, the ratio of observed CO(2-1) to CO(1-0) intensity ( $R_{21/10}$ ) varies systematically with star formation rate and gas surface density in M51 (Koda *et al.* 2012). CO(2-1) likely traces the denser and/or warmer gas, rather than the bulk molecular gas in galaxies.

## References

- Bigiel, F. *et al.* 2008, *AJ*, 136, 2846  
Kennicutt, Jr., R. C., 1998, *ARAA*, 36, 189  
Koda, J. *et al.* 2012, accepted *ApJ*  
Leroy, A. K. *et al.* 2008, *AJ*, 136, 2782  
Schmidt, M., 1959, *ApJ*, 129, 243  
Tan, J. C., 2010, *ApJ*, 710, L88